

Reptilia, Squamata, Serpentes, *Lygophis paucidens* Hoge, 1952: First records for Paraguay

Pier Cacciali^{1,2*}, Paul Smith^{1,3}, Anna Källberg¹, Helen Pheasey¹ and Karina Atkinson¹

¹ Para La Tierra, Reserva Natural Laguna Blanca, Santa Rosa del Aguaray, Departamento San Pedro, Paraguay.

² Consejo Nacional de Ciencia y Tecnología – Instituto de Investigación Biológica del Paraguay. Del Escudo 1697, Asunción, Paraguay.

³ Fauna Paraguay. Encarnación, Paraguay.

*Corresponding author. E-mail: pier_cacciali@yahoo.com

ABSTRACT: The first Paraguayan records of the poorly known xenodontine snake *Lygophis paucidens* are reported. Two male specimens were collected in a mosaic habitat of *Campo Sucio* and *Cerrado sensu stricto* in the Reserva Natural Laguna Blanca, Departamento San Pedro. This is the first record of this species outside of Brazil.

Lygophis paucidens Hoge, 1952 is a poorly known xenodontine snake associated with Caatinga/Cerrado habitats in east-central Brazil (Michaud and Dixon 1987; Nogueira 2001; Tipton 2005; Valdujo *et al.* 2009). The species was originally described from Mato Verde, Mato Grosso, Brazil (Hoge 1952), but despite very few specimens the distribution appears to be wide, with further documented records from the Brazilian States of Piauí, Pernambuco, Paraíba, Goiás, and Distrito Federal (Tipton 2005; Valdujo *et al.* 2009). However records from Cerrado are few and by 2001 the species was known from just 10 specimens in six Brazilian Cerrado localities in the states of Goiás and Distrito Federal (Nogueira 2001; Tipton 2005), with an additional record from Goiás later published by Valdujo *et al.* (2009).

Lygophis paucidens can be distinguished from related taxa such as *L. dilepis* by the presence of dark longitudinal lines only on the anterior part of the body, which abruptly fade a short distance behind the head. Published scale counts for this species are: ventrals 166–174; subcaudals 62–72; and 17-17-15 dorsal scale rows (Lema 1989). Because of its rarity, little is known about the biology of *L. paucidens*. The only published ecological information on the species is a comment by Michaud and Dixon (1989) that *L. paucidens* feeds on small teiid lizards of the genus *Cnemidophorus*, which are widespread and generally common throughout Caatinga and Cerrado (Colli *et al.* 2002).

In this note we report on two specimens of *L. paucidens* from Cerrado habitat at Reserva Natural Laguna Blanca (RNLB), Departamento San Pedro, Paraguay (Figure 1) (23°47'56"S, 56°17'32"W), which represent the first record for the country and the first report of this species outside Brazil.

The RNLB is an 804 hectare reserve consisting of over 400 hectares of near pristine Cerrado, a patch of degraded Atlantic Forest and areas of transitional semi-deciduous, semi-humid forest. The four main Cerrado ecotypes are present at RNLB and grow on a predominately sandy substrate (Eiten 1972, 1978). The RNLB was declared a Natural Reserve on 3 February 2010 for a limited period of just five years (Decree 3893 under Article 26 of the

Protected Areas Law 352/94).

The following specimens were collected by fieldworkers from Estación Biológica Para La Tierra (PLT), based at RNLB and are deposited in Para La Tierra Biological Station collection (CZPLT) located at the reserve. Both specimens were collected in a mosaic of *Campo Sucio* and *sensu stricto* Cerrado vegetation types within the reserve (*sensu* Eiten, 1972, 1978).

The specimen CZPLT-H-122 (March 15, 2011) is male, collected in a pitfall trap (Figures 2 and 3). Measurements: total length 401.5 mm, snout-vent length 350.0 mm, dorsal scale rows 17-17-15, ventrals 165, subcaudals 66. The




FIGURE 1. Map showing the locality record of *Lygophis paucidens* in Paraguay.

ventral scale count of this specimen extends the lower limit of the known range for the species.

The specimen CZPLT-H-144 (May 2, 2011) is also male and was collected under a metal Sherman trap on a cold morning. Measurements: total length 390.0 mm, snout-vent length 290.5 mm, dorsal scale rows 17-17-15, ventrals 167, subcaudals 63.

The records reported here represent a range extension of more than 320 Km SW from the closest published locality Assentamento Nascentes do Araguaia, Goiás State (Valdujo *et al.* 2009) and confirm a wider distribution of *L. paucidens* throughout the Cerrado region.

Though small, the importance of the RNLB for conservation in Paraguay should not be underestimated. With over 400 hectares of globally threatened, pristine Cerrado habitat, plus an artesian lake arising from the Guarani Aquifer, the reserve is of both national and international importance. It was declared an Important Bird Area by Birdlife International (IBA PY021- Guyra

Paraguay 2008) due to the extraordinarily high number of globally (11) and nationally (47) threatened bird species occurring within the reserve. Indications from ongoing herpetological inventories suggest that it is also an area of national and international importance for the conservation of reptiles and amphibians, given that despite its small size it boasts the highest amphibian and reptile species lists of any protected area in Paraguay (Smith *et al.* 2012, Para La Tierra unpublished data). Consolidating the long term conservation of the RNLB should be a national conservation priority.

ACKNOWLEDGMENTS: Thanks to Francisco L. Franco for revising the manuscript. Dr Norman Scott has been a constant source of knowledge and inspiration to all of us and deserves special thanks for his unwavering support of the work done by Para La Tierra. We thank SEAM for issuing the relevant permits and for their recognition of the importance of RNLB by declaring it a Reserva Privada. Particular thanks to Malvina Duarte the owner of Laguna Blanca for her foresight and support of PLT, without which the fieldwork would not have been possible. Thanks to all the volunteers and staff at Para La Tierra Biological Station for all their efforts in documenting the understudied fauna of the Paraguayan cerrado. Specimens were collected under permit 03/11 issued by the Secretaría del Ambiente.

LITERATURE CITED

- Colli, G. R., P. Bastos and A. F. Araujo. 2002. The character and dynamics of the Cerrado herpetofauna; p. 223-241 In P. S. Oliveira and R. J. Marquis (eds.) *The cerrados of Brazil: ecology and natural history of a neotropical savanna*, New York: Columbia University Press.
- Nogueira, C. N. C. 2001. New records of Squamate reptiles in central Brazil Cerrado II: Brasília region. *Herpetological Review* 32: 285-287.
- Eiten, G. 1972. The Cerrado vegetation of Brazil. *Botanical Review* 38: 201-341.
- Eiten, G. 1978. Delimitation of the Cerrado concept. *Vegetatio* 36: 169-178.
- Guyra Paraguay. 2008. Áreas importantes para la conservación de las aves del Paraguay. Asunción: Asociación Guyra Paraguay. 473 p.
- Hoge, A. 1952. Notas hepitológicas. Primeira contribuição ao conhecimento dos ofídios do Brasil central. *Memorias do Instituto Butantan* 24: 179-214.
- Lema, T. de. 1989. Serpentes do complexo *Liophis lineatus* (Linnaeus, 1758) no Brasil nordeste (Serpentes, Colubridae: Colubrinae). *Acta Biologica Leopoldensia* 11: 251-271.
- Michaud, E. and J. Dixon. 1987. Taxonomic revision of the *Liophis lineatus* complex (Reptilia: Colubridae) of Central and South America. *Contributions in Biology and Geology* 71: 1-26.
- Michaud, E. and J. Dixon. 1989. Prey items of 20 species of the neotropical colubrid snake genus *Liophis*. *Herpetological Review* 20: 39-41.
- Smith, P., P. Cacciali, K. Atkinson, H. Pheasey and M. Motte. 2012. New distributional records of amphibians for Departamento San Pedro, Paraguay (Amphibia). *Check List* 8(5): 903-907.
- Tipton, B. 2005. Snakes of the Americas, checklist and lexicon. Florida: Krieger Publishing Company. 492 p.
- Valdujo, P. H., C. de Campos Nogueira, L. Baumgarten, F. H. Guimaraes Rodrigues, R. Albuquerque Brandao, A. Eterovic, M. Barroso Ramos-Neto and O. A. V. Marques. 2009. Squamate reptiles from Parque Nacional das Emas and surroundings, Cerrado of Central Brazil. *Check List* 5: 405-417.

RECEIVED: August 2012

ACCEPTED: December 2012

PUBLISHED ONLINE: March 2013

EDITORIAL RESPONSIBILITY: Ross MacCulloch



FIGURE 2. Specimen of *Lygophis paucidens* (CZPLT-H-122) found at Reserva Natural Laguna Blanca, in San Pedro Department, Paraguay.



FIGURE 3. Detail of the head of *L. paucidens* (CZPLT-H-122).